

RECEIVED
CENTRAL FAX CENTER

MAR 06 2007

Atty Dkt. No.: 10031551-1
USSN: 10/813,467**AMENDMENTS TO THE CLAIMS**

Please incorporate the following amendments to the subject application.

In the Claims:

1-27 (Cancelled)

28. (Currently Amended) A method for ~~producing~~ synthesizing an addressable array of oligonucleotides on a substrate, comprising:
contacting a substrate comprising a plurality of addressable features ~~comprising a surface~~ ~~substrate~~ attached blocked nucleoside monomer blocked with a blocking group, or a ~~substrate surface~~ attached polymer blocked with a blocking group, with a deblocking fluid to remove the blocking group, thereby generating an unblocked attached nucleoside monomer or polymer;
displacing the deblocking fluid from the substrate surface comprising the attached unblocked nucleoside monomer or polymer with a purging fluid; and
reacting the attached unblocked nucleoside monomer or polymer with another blocked nucleoside monomer to produce said addressable array of oligonucleotides.

29. (Previously Presented) The method of claim 28, wherein a blocked nucleoside monomer is attached to the substrate by contacting the substrate with a fluid comprising a blocked nucleoside monomer at a location on the substrate that comprises hydroxyl functional groups.

30. (Previously Presented) The method of claim 28, wherein steps of the method are repeated a plurality of times.

31. (Previously Presented) The method of claim 28, wherein the substrate comprises a surface of a planar support.

BEST AVAILABLE COPY

Atty Dkt. No.: 10031551-1
USSN: 10/813,467

32. (Previously Presented) The method of claim 28, wherein the displacing step causes minimal mixing of deblocking and purging fluids.
33. (Currently Amended) The method of claim ~~[[28]]~~ 56, wherein the substrate comprises a surface of a support containable within a flow cell.
34. (Currently Amended) The method of claim 28, wherein the substrate comprises a nascent surface of a planar support.
35. (Previously Presented) The method of claim 28, wherein the purging fluid has a density that is different from the deblocking fluid.
36. (Currently Amended) The method of claim 28, wherein the purging fluid and the deblocking fluid have a density difference, expressed as the Atwood number (A), of at least ~~about~~ 0.01.
37. (Previously Presented) The method according to claim 28, wherein the purging fluid has a density that is higher than the density of the deblocking fluid.
38. (Previously Presented) The method according to claim 28, wherein the purging fluid has a density that is lower than the density of the deblocking fluid.
39. (Previously Presented) The method according to claim 28, wherein the purging fluid is an organic fluid.
40. (Previously Presented) The method according to claim 28, wherein the purging fluid comprises an oxidizing agent.
41. (Previously Presented) The method according to claim 28, wherein the purging fluid comprises a wash fluid.
42. (Previously Presented) The method according to claim 41, wherein the wash fluid is an organic fluid.

Atty Dkt. No.: 10031551-1
USSN: 10/813,467

43. (Previously Presented) The method according to claim 41, wherein the wash fluid is acetonitrile.

44. (Previously Presented) The method according to claim 28, wherein deblocking fluid is displaced from the surface with a purging fluid by flowing the purging fluid across the surface in a manner sufficient to produce a stratified fluid interface that moves across the surface.

45. (Previously Presented) The method according to claim 44, wherein the purging fluid is flowed across the surface at a rate ranging from about 1 cm/s to about 20 cm/s.

46. (Previously Presented) The method of claim 28, wherein the purging fluid limits the efficiency of deblocking by the deblocking fluid.

47. (Previously Presented) The method of claim 29, wherein the hydroxyl functional groups are provided by 5'-OH groups of nucleoside monomers or polymers attached to the substrate.

48. (Previously Presented) The method of claim 28, wherein the step of displacing occurs in a flow cell.

49. (Previously Presented) The method of claim 29, wherein the blocked nucleoside monomer is deposited at the location by pulse jetting.

50. (Previously Presented) The method of claim 28, wherein the blocking group comprises an acid labile blocking group and wherein the deblocking fluid comprises an acid.

51. (Previously Presented) The method of claim 33, wherein the substrate is contained within a chamber of a flow cell when contacted with deblocking fluid and wherein the chamber comprises at least one fluid inlet and at least one fluid outlet.

Atty Dkt. No.: 10031551-1
USSN: 10/813,467

52. (Previously Presented) The method of claim 51, wherein the flow cell is oriented in an at least partially vertical position.

53. (Previously Presented) The method of claim 44, wherein a pressure gradient is used to produce the stratified interface.

54. (Currently Amended) The method of claim 28, wherein the deblocking fluid comprises an organic solvent that has a vapor pressure that is less than **about** 13 Kpa at 0 °C and 1 ATM.

55. (Previously Presented) The method of claim 28, further comprising contacting the substrate comprising the attached blocked nucleoside monomer or polymer with an oxidation fluid prior to contacting with the deblocking fluid.

56. (Currently Amended) A method of producing an addressable array comprising a substrate of and at least two oligonucleotides bonded to different locations on a surface of the said substrate, comprising:

contacting nucleoside monomers blocked with a blocking group to at least a first location and a second different location of a said substrate surface displaying functional groups, under conditions sufficient for the blocked nucleoside monomers to bond to the surface in first and second locations to produce a substrate surface displaying bound blocked monomers;

contacting the surface displaying bound blocked monomers with a deblocking fluid to remove the blocking group, thereby generating unblocked nucleoside monomers at the first and second locations;

displacing the deblocking fluid from the surface comprising the bound unblocked monomers at the first and second locations with a purging fluid;

reacting the attached unblocked nucleoside monomers at the first and second locations with another blocked nucleoside monomer to produce said addressable array.

BEST AVAILABLE COPY

Atty Dkt. No.: 10031551-1
USSN: 10/813,467

57. (Previously Presented) The method of claim 56, wherein the at least two oligonucleotides comprise the same sequence composition.

58. (Previously Presented) The method of claim 56, wherein the at least two oligonucleotides comprise different sequence compositions.

59. (Previously Presented) The method of claim 56, further comprising contacting the bonded blocked monomers with an oxidation fluid prior to contacting the surface with the deblocking solution.

60. (Cancelled)

New Claim:

61. (New) The method according to claim 28, wherein said substrate is planar.